

List of Tables

2-1	Size and estimated age of maturity of California abalones	2-2
2-2	Biological information summary	2-3
2-3	Growth parameters of California abalones summary	2-5
2-4	Red abalone population survey at Van Damme, Fort Ross and Salt Point State Parks, 1986-2000	2-18
6-1	Interim and long-term abundance and depleted areas as criteria for recovery	6-4
6-2	List of recovery activities with field time required	6-13
6-3	Key locations for recovery of red abalone in southern California	6-18
6-4	Key locations for recovery of pink abalone in southern California	6-20
6-5	Key locations for recovery of green abalone in southern California	6-22
6-6	Key locations for recovery of black abalone in southern California	6-24
6-7	Key locations for recovery of white abalone in southern California	6-26
6-8	Key locations for recovery of pinto and flat abalones in southern California	6-27
6-9	Estimates for the amount of time for recovery of five species of abalone in southern California	6-28
7-1	Abalone survey summaries, 1999-2000	7-4
7-2	Total Allowable Catch adjustment decision table using established criteria	7-6
7-3	Projected changes (percent) in the baseline TAC (400,000 abalone) with various combinations of daily and annual limits	7-8
7-4	Site closure decision table using established criteria	7-11
9-1	Timeline of recovery activities for 2003 to 2009	9-5
9-2	Timeline of interim management activities for 2003 to 2009	9-6
9-3	Summary of costs and funding sources for implementation averaged over 2003 to 2009	9-7